

Signature of the $\Lambda(1405)$ resonance in neutron spectra from the $K^- + d$ reaction

J. Révai¹

Bogoliubov Laboratory of Theoretical Physics, JINR, 141980 Dubna, Russia

Wigner Research Center for Physics, 1525 Budapest, POB 49, Hungary

Abstract

Neutron spectra from the reaction $K^- + d \rightarrow \pi + \Sigma + n$ were calculated in the energy range $E_{K^-}^{cm} = 0 - 50 \text{ MeV}$ using coupled channel Faddeev equations for the description of the $\bar{K}NN - \pi\Sigma N$ three-body system. The aim was to trace the signature of the $\Lambda(1405)$ resonance in the spectra. We found, that while in the direct spectra kinematic effects mask completely the peak corresponding to the resonance, the deviation spectrum method [1] is able to eliminate kinematics and produce clear evidence of the resonance. Four different phenomenological $\bar{K}N - \pi\Sigma$ interactions were used in order to study the effect of $\Lambda(1405)$ pole position on the neutron spectra.

Keywords: kaon-baryon interactions, Faddeev equations, kaon induced reactions and scattering, nuclear few-body systems

1. Introduction

The $\Lambda(1405)$ resonance plays a central role in low-energy kaon-nuclear physics. Being a manifestation of the assumed attraction between negative antikaons and nucleons its observability and properties are crucial for the possible existence of antikaonic nuclear clusters. Its origin and structure, mainly its one- or two-pole nature, the position and widths of these poles, are subject to vivid discussions among the representatives of different opinions, in particular, chiral perturbation theory versus phenomenology. For a review of the "state of art" see e.g. [1, 2] and references therein. From experimental

¹email: revai.janos@wigner.mta.hu

point of view, the clarification of this problem is hindered by the fact, that this state can not be reached in two-body reactions involving stable particles, it can be observed only in processes having 3 or more particles in final states. The simplest of these is the reaction

$$\begin{array}{ccc}
 & \nearrow K^- + d \text{ or } K^- + p + n & \\
 K^- + d & \rightarrow (\pi + \Sigma)_{I=0,1} + n & (1) \\
 & \searrow (\pi + \Sigma)_{I=1} + p &
 \end{array}$$

which has the advantage, that its dynamics can be treated exactly in the framework of the coupled particle-channels Faddeev approach. This formalism has been applied earlier for this 3-body system mainly for calculation of 3-body quasi-bound states [3, 4], or K^-d scattering length [5, 6, 7]. The break-up reaction (1) was studied in the early papers [8, 9] (of course, together with values for the K^-d scattering length, too) but with the main emphasis on the possible signal of a resonance in the $\Lambda N - \Sigma N$ system, close to the ΣN threshold. Probably at that time the $\Lambda(1405)$ topic was not as hot as nowadays.

The only available experimental data on neutron spectra from the reaction (1) are those of Tan [10] from a bubble chamber experiment. What he gets is a large peak near the origin with no distinguishable structure in it. It corresponds to our "direct" spectrum $P(E_n)$ (see later).

The recent papers [11, 12] and [2] are devoted to the possible observation of the $\Lambda(1405)$ in the reaction (1). In these papers the dynamics is treated in single- plus double-scattering approximation, which might be justified at higher incident kaon energies. Our calculation is done for low-energy kaons, in the range $E_{K^-}^{cm} = 0 - 50 \text{ MeV}$, having in mind stopped or slowed down kaons.

The paper is organized as follows: in Sect. II. we give a brief description of the applied formalism, Sect. III. contains the details and the input of the calculation, our results and their discussion are presented in Sect. IV., while our conclusions are in Sect. V.

2. Formulation of the problem

The calculation is based on the coupled-channels AGS-Faddeev treatment of the $\bar{K}NN - \pi\Sigma N$ three-body system. The details of this approach have been already described in detail in several papers [3, 4, 5], here we shall

recall them only briefly, mainly to introduce the notations. The operator AGS equations for the transition operators U_{i1} read

$$U_{i1} = (1 - \delta_{ij})G_0^{-1} + \sum_{j \neq i} T_j G_0 U_{j1}, \quad (2)$$

where $i, j = 1, 2, 3$ are the usual pair-spectator indices,

$$T_j = V_j + V_j G_0 T_j$$

are the two-particle T -operators and $G_0(z) = (z - H_0)^{-1}$ is the free Green-operator.

The configuration space $|\mathbf{x}_i \mathbf{y}_i \nu_i\rangle$ in which these operators act, apart from the usual Jacobi momentum variables $\mathbf{x}_i \mathbf{y}_i$ contain a discrete index $\nu_i = (\alpha, \sigma_i)$, which is a combination of the particle composition index α

$$\alpha = \{1, 2, 3\} = \{\bar{K} N_1 N_2, \pi \Sigma_1 N_2, \pi N_1 \Sigma_2\}$$

and an isospin label $\sigma_i = (I_i I)$ corresponding to pair isospin I_i of the particle pair i and total isospin I :

$$\sigma_i \sim \left[[t_j t_k]^{I_i} t_i \right]^I$$

This choice of the isospin labels corresponds to the "isospin representation", which is useful when isospin conserving pair interactions are used. Another possibility is the equivalent "charge state" or "particle" representation, characterized by the 3rd component of the particle isospins:

$$\sigma_0 \sim \{t_{1z}, t_{2z}, t_{3z}\} \quad \text{or for } \alpha = 1 \quad \sigma_0 \sim \{\bar{K}^0 n_1 n_2, K^- p_1 n_2, K^- n_1, p_2\}$$

These representations can be transformed into each other with the help of orthogonal matrices

$$B_{ij}^\alpha = \langle \alpha \sigma_i | \alpha \sigma_j \rangle \quad i, j = 0, 1, 2, 3 \quad (3)$$

composed from 6j symbols for $i, j = 1, 2, 3$ and from Clebsch-Gordan coefficients for $i = 0, j = 1, 2, 3$. Using isospin conserving separable interactions of the form

$$V_{\nu_i \nu'_i}^i = \delta_{I_i I'_i} \delta_{II'} |g_{\nu_i}^{I_i}\rangle \lambda_{\nu_i \nu'_i}^{I_i} \langle g_{\nu'_i}^{I_i}| \quad (4)$$

the $T_i(z)$ operators can be written as

$$T_i(z) = \sum_{\nu_i \nu'_i} |g_{\nu_i}^{I_i}\rangle \tau_{\nu_i \nu'_i}(z) \langle g_{\nu'_i}^{I'_i}| \quad (5)$$

with $\tau_{\nu_i \nu'_i}(z)$ being the usual (c-number) matrix, defined as:

$$(\tau_{\nu_i \nu'_i}(z))^{-1} = (\lambda_{\nu_i \nu'_i}^{I_i})^{-1} - \langle g_{\nu_i}^{I_i} | G_0(z) | g_{\nu'_i}^{I'_i} \rangle \quad (6)$$

The matrix indices ν_i in Eq.(6) consist of the particle space label α and the isospin label $\sigma_i = (I_i I)$. Due to isospin conservation of our interactions, the coupling constant matrix λ is diagonal in $(I_i I)$, while for particle pairs i capable to change their identity ($\bar{K}N \leftrightarrow \pi\Sigma$) it has non-diagonal elements in the particle labels $(\alpha\alpha')$. As for the matrix elements of G_0 , it does not change particle identities, thus it is diagonal in α , and if we take averaged masses for particles within an isospin multiplet, it is also diagonal in pair- and total isospin indices $(I_i I)$. However, if physical (unequal) masses are used, G_0 will be diagonal only in "particle" representation, while in the σ_i "isospin" representation it will acquire non-diagonal elements both in I_i and I , proportional to the mass differences.

The equations (2) for the transition operators take the form

$$U_{i1} = (1 - \delta_{ij})G_0^{-1} + \sum_{j \neq i} \sum_{\nu'_j} |g_{\nu_j}\rangle \tau_{\nu_j, \nu'_j} \langle X_{\nu'_j}^j| \quad \text{with} \quad \langle X_{\nu'_j}^j| = \langle g_{\nu'_j}^{I_j} | G_0 U_{j1} \quad (7)$$

Introducing the functions $X_{\nu_j}^j(\mathbf{y}_j) = \langle X_{\nu_j}^j | \Phi_0 \rangle$, where $\mathbf{y}_i$ is the momentum of the spectator particle, corresponding to the pair j and $|\Phi_0\rangle = |\varphi_d P_K\rangle$ is the initial state with the deuteron wave function $|\varphi_d\rangle$ and P_K - the momentum of the incident kaon, we get the set of integral equations:

$$X_{\nu_i}^i(\mathbf{y}_j) = (1 - \delta_{i1}) \langle g_{\nu_i}^{I_i} | \Phi_0 \rangle + \sum_{j \neq i} \sum_{\nu'_j} \int Z_{\nu_i \nu_j}(\mathbf{y}_i, \mathbf{y}_j) \tau_{\nu_j \nu'_j}(z - y_j^2) X_{\nu'_j}^j(\mathbf{y}_j) d\mathbf{y}_j \quad (8)$$

with the kernel

$$Z_{\nu_i \nu_j}(\mathbf{y}_i, \mathbf{y}_j) = \langle g_{\nu_i}^{I_i} | G_0(z) | g_{\nu_j}^{I_j} \rangle. \quad (9)$$

The size of the system (9) can be reduced by introducing symmetric (antisymmetric) combinations of X-functions, with respect to interchange of baryon

numbering. The baryon spins do not enter explicitly in this formalism, therefore the total baryon spin S remains unchanged in the process (is a conserved quantum number). For a given S value the total antisymmetry required by the Pauli principle has to be ensured by the space-isospin part. Thus for $S = 0$ (K^-pp system) we have to work with the symmetric combinations of X -s, while for $S = 1$ (our K^-d system) the antisymmetric combinations are needed. As a result, the labeling of the unknown functions $\nu_i = (\alpha\sigma_i)$ is changed to $\mu_a = (a, \sigma_a)$, where a denotes a pair of interacting particles, irrespectively to which original particle composition channel they belonged and σ_a denotes the corresponding isospin values. Thus we are left with $X_{a\sigma_a}(\mathbf{y}_a)$ and a can take the values $\bar{K}N, NN, \Sigma N$ and $\pi\Sigma$ (πN is missing, since we neglected the πN interaction, see next section).

The break-up transition operator U_{01} can be expressed in terms of the U_{i1} -s as:

$$U_{01} = \frac{1}{2}(U_{11} + U_{21} + U_{31})$$

and the break-up amplitude reads

$$A_{BU} = \langle \Phi_f | U_{01} | \Phi_0 \rangle. \quad (10)$$

For the reaction under consideration the properly antisymmetrized final state is

$$|\Phi_f\rangle = |\mathbf{x}_{\pi\Sigma}, \mathbf{y}_N; \sigma_{\pi\Sigma}\rangle = \frac{1}{\sqrt{2}}(|\mathbf{x}_{\pi\Sigma_1}, \mathbf{y}_{N_2}; \sigma_{\pi\Sigma_1}\rangle - |\mathbf{x}_{\pi\Sigma_2}, \mathbf{y}_{N_1}; \sigma_{\pi\Sigma_2}\rangle)$$

The break-up amplitude can be expressed in terms of the X -functions as

$$\begin{aligned} A_{BU}(\mathbf{x}_{\pi\Sigma}, \mathbf{y}_N; \sigma_{\pi\Sigma}) = & \\ - g_{\pi\Sigma}(\mathbf{x}_{\pi\Sigma}) & \left[\tau_{\pi\Sigma, \bar{K}N}(z - y_N^2) X_{\bar{K}N}(\mathbf{y}_N) + \tau_{\pi\Sigma, \pi\Sigma}(z - y_N^2) X_{\pi\Sigma}(\mathbf{y}_N) \right] \\ - B_{31}^2 g_{\Sigma N}(u\mathbf{y}_N + v\mathbf{x}_{\pi\Sigma}) & \tau_{\Sigma N, \Sigma N}(z - |\mathbf{x}_{\pi\Sigma} - w\mathbf{y}_N|^2) X_{\Sigma N}(\mathbf{x}_{\pi\Sigma} - w\mathbf{y}_N), \end{aligned} \quad (11)$$

where B_{31}^2 is an isospin recoupling matrix (see Eq.(3)), u, v and w are mass coefficients of the transformation between Jacobi momentum sets. In Eq.(11) we omitted the isospin labels, the quantities are vectors (matrices) in isospin space. The on-shell amplitude for a given neutron energy E_n depends on E_n, t and the isospin labels $\sigma_{\pi\Sigma}$:

$$A(E_n, t, \sigma_{\pi\Sigma}) = A_{BU}(\mathbf{x}_{\pi\Sigma}, \mathbf{y}_N; \sigma_{\pi\Sigma})$$

with

$$|\mathbf{y}_n| = \sqrt{2E_n\mu_{N,\pi\Sigma}}; \quad |\mathbf{x}_{\pi\Sigma}| = \sqrt{2(E_{\pi\Sigma N} - E_n)\mu_{\pi\Sigma}}; \quad t = \cos(\mathbf{x}_{\pi\Sigma}, \mathbf{y}_n)$$

The physically observable final state corresponds to a certain particle composition, not to a definite isospin state, therefore the amplitude has to be transformed into the σ_0 representation, using the suitable B matrix of Eq.(3):

$$A(E_n, t, \sigma_0) = \sum_{\sigma_{\pi\Sigma}} (B_{03}^2)_{\sigma_0, \sigma_{\pi\Sigma}} A(E_n, t, \sigma_{\pi\Sigma}),$$

where σ_0 can be $\{\pi^+\Sigma^-n, \pi^0\Sigma^0n, \pi^-\Sigma^+n\}$. The neutron spectrum is proportional to the differential cross section

$$P(E_n, t, \sigma_0) \sim \frac{d\sigma}{d\Omega_{\mathbf{x}_{\pi\Sigma}} d\Omega_{\mathbf{y}_N} dE_n} = (2\pi)^4 \mu_{\pi\Sigma} \mu_{N,\pi\Sigma} \mu_{K,NN} \frac{x_{\pi\Sigma} y_N}{P_K} |A(E_n, t, \sigma_0)|^2$$

The inclusive neutron spectrum (when no other particles are detected) is given by

$$P(E_n) = \sum_{\sigma_0} \int_{-1}^1 dt P(E_n, t, \sigma_0) \quad (12)$$

The above considerations refer to the neutrons emerging from the reaction $K^- + d \rightarrow \pi + \Sigma + n$; when the energy of the incident kaon exceeds the deuteron binding energy, neutrons are also emitted from the reaction $K^- + d \rightarrow K^- + p + n$. Their spectrum can be deduced in a similar way to Eq.(11) from the $X_{\bar{K}N}$ and X_{NN} functions. The allowed energy range for neutrons from the first reaction is $(0, E_K + E_d + \Delta)$, while for the second it is $(0, E_K + E_d)$, where Δ is the difference of the $\bar{K}N$ and $\pi\Sigma$ threshold energies.

3. Details of the calculation and the input

The main purpose of the present work is to study the possible signature(s) of the $\Lambda(1405)$ resonance in the neutron spectra from the reaction (1). In our calculation we used the two-body interactions of [5] and [6], which are adjusted for our three-body model. They are s -wave, separable isospin dependent and isospin conserving interactions with Yamagouchi type form-factors.

In particular, for the two-channel $\bar{K}N - \pi\Sigma$ interaction we used two variants, both having a one and a two pole version for the $\Lambda(1405)$. They

both reproduce all available experimental data on the low-energy $\bar{K}N$ system, the first one is fitted to the KEK data on the kaonic hydrogen $1s$ level shift, while the second one reproduces the most recent SIDDHARTA data. Their pole positions relevant for the $\Lambda(1405)$ are shown in Table 1 (the negative real parts correspond to distances from the $\bar{K}N$ threshold).

Table 1: Pole positions of the $\bar{K}N - \pi\Sigma$ potentials (in MeV)

	KEK	SIDDHARTA
1-pole	-23.6-35.6 i (1411.0-35.6 i)	-6.4-46.8 i (1428.1-46.8 i)
2-pole	-22.2-36.3 i (1412.4-36.3 i)	-14.8-57.2 i (1419.7-57.2 i)

The numbers in Table 1 differ slightly from those given in the original papers [5, 6]. The reason is, that the above ones were calculated with averaged masses and without Coulomb interaction - as they appear in most of the 3-body calculations, - while the fitting to the experimental data was performed with physical masses and Coulomb interaction. Since the main aim of the present work is to study the appearance of subthreshold resonances of different type in a 3-body reaction, we kept both interactions, not only the more advanced one.

The triplet NN interaction is a two-term one to account for the short range repulsion, with form-factors fitted to reproduce the deuteron and s -wave phase shifts. The $S = 1$ ΣN interaction in the $I = 1/2$ isospin state is complex, since it was deduced from a two-channel $\Sigma N - \Lambda N$ interaction, while for $I = 3/2$ it is real. The πN interaction was neglected in our calculation due to its weak s -wave part.

The total angular momentum was restricted to $L = 0$ since we believe, that for our s -wave interactions the essential dynamics can be traced in spite of this limitation. Keeping the interactions s -wave, the extension to higher partial waves is straightforward, unlike the case of inclusion of $p, d, \dots$ -wave interactions.

We considered incident kaon energies in the interval $E_{K^-}^{cm} = 0 - 50$ MeV ($P_K^{LAB} \sim 0 - 250$ MeV/c). The system of integral equations (9) in the case of physical masses ($I = 1/2$ and $I = 3/2$ mixed) consists of 12 equations, while for averaged masses - of 8.

As a numerical method we used expansion of the unknown functions on a cubic spline basis, for the distribution of nodes and collocation points the

prescription of [13] was used with a slight modification to allow nonsymmetric intervals and distributions on the two sides of the break-up singularity. Complete convergence of the results was achieved for ~ 20 nodes in the non-break-up channels, while for the break-up channels $\sim 30 - 35$ nodes were necessary. Apart from the lower dimensionality of the matrices to be inverted, the use of spline expansion is especially advantageous when break-up amplitudes are calculated, since no interpolation of the solutions is needed.

4. Results and discussion

We start the presentation of our results by a "by-product": the effect of the physical versus averaged masses of the $\bar{K}^0$ and K^- mesons on the K^-d scattering length. The inclusion of the possibility of isospin mixing due to this mass difference allowed us to extend the results of [5, 6] in this respect. Our results for the two potential versions are shown in Table 2. The results for averaged masses coincide with those of [5, 6], while for physical masses they differ by a few per cent, mainly in the real part. At present level of accuracy of available information - both theoretical and experimental - on the $\bar{K}N$ interaction and $\bar{K}$ -nuclear clusters this difference does not seem to be essential. However, once it might become useful to have some numerically reliable information on the order of magnitude of this effect.

Table 2: K^-d scattering lengths of the $\bar{K}N - \pi\Sigma$ potentials for physical and averaged masses of K^- and $\bar{K}^0$ (in fm).

	KEK		SIDDHARTA	
	averaged	physical	averaged	physical
1-pole	-1.49+0.97 i	-1.52+0.98 i	-1.47+1.22 i	-1.50+1.23 i
2-pole	-1.57+1.10 i	-1.60+1.12 i	-1.50+1.23 i	-1.54+1.24 i

On the other hand, when calculating neutron spectra our interest was focused on qualitative signals of the $\Lambda(1405)$ in the line shapes, therefore we used averaged masses, what simplified the numerical work to some extent.

We have calculated the inclusive neutron spectra $P(E_n)$ (12) for different incident kaon energies, both below and above the deuteron break-up threshold. For kaon incident energies $E_{K^-}^{cm} = 1, 20, 50 MeV$ the results are displayed

in Figs. 1, 2 and 3, respectively, (upper left graphs)².




Figure 1: Neutron spectra for kaon incident energy $E_{K^-}^{cm} = 1 \text{ MeV}$. Upper left graph - direct neutron spectra $P(E_n)$, eq.(12); lower left graph - deviation spectrum $P_{DEV}(E_n)$, eq.(17); upper right graph - single scattering deviation spectrum $P_{DEV}^{sing}(E_n)$, eq. (17); lower right graph - "original" $\Lambda(1405)$ as $\pi^0 \Sigma^0$ elastic cross section.

The overall shape of the spectra is a strong peak near the origin with no signal of the $\Lambda(1405)$ resonance. The direct $P(E_n)$ spectra are practically

²Since our main concern is the possible trace of the $\Lambda(1405)$ in the line shapes of the calculated spectra, the arbitrary units on the Y-axes of our graphs are chosen to optimize visibility.

indistinguishable for the four considered $\bar{K}N - \pi\Sigma$ potentials. For kaon energies above the deuteron binding energy there are two modifications: the neutron spectrum from the $K^- + d \rightarrow \pi + \Sigma + n$ channel show a cusp at neutron energies $E_n = E_{th}$ when the $\bar{K}N$ system is at its threshold, and additional neutrons show up from the $K^- + d \rightarrow K^- + p + n$ reaction in a form of a structureless bump between $E_n = 0$ and $E_n = E_{th}$ (on the graphs it is scaled




Figure 2: Neutron spectra for kaon incident energies $E_K^{cm} = 20 \text{ MeV}$ The four graphs are the same as in Fig.1.

down to allow to draw it on the same plot with the other neutrons). The reason, why the $\Lambda(1405)$ is not seen in these spectra is essentially kinematical: the neutron energy in the resonance region should exceed the incident energy

of the kaon by $10 - 20 - 30 \text{ MeV}$, depending on the resonance position, while in the deuteron the neutron energy (momentum) distribution is dominated by the low energy part.




Figure 3: Neutron spectra for kaon incident energies $E_{K^-}^{cm} = 50 \text{ MeV}$ The four graphs are the same as in Fig.1.

In order to eliminate this kinematical "inconvenience" Esmaili, Akaishi and Yamazaki (EAY) [1] propose to consider instead of $P(E_n)$ the deviation spectrum:

$$P_{DEV} = \frac{P(E_n)}{P_{nonres}(E_n)} \quad (13)$$

In Eq.(13) $P_{nonres}(E_n)$ is a non-resonant background spectrum, containing

the kinematics of the reaction. Let's see, how this idea can be realized in our case.

Considering the zero-order iteration of our (symmetrized) system of integral equations (9), the only non-vanishing X will be the inhomogeneous term:

$$X_{\bar{K}N}(y_N) = \langle g_{\bar{K}N} | \Phi_0 \rangle = \langle g_{\bar{K}N} | \varphi_d P_K \rangle. \quad (14)$$

This ansatz is usually called single scattering approximation. Substituting eq.(14) into eq.(11) we get the corresponding break-up amplitude:

$$\begin{aligned} A_{BU}^{sing}(x_{\pi\Sigma}, y_N; \sigma_{\pi\Sigma}) &= g(x_{\pi\Sigma}) \tau_{\pi\Sigma, \bar{K}N}(z - y_N^2) \langle g_{\bar{K}N} | \varphi_d P_K \rangle \\ &= \langle x_{\pi\Sigma}, y_N | T_{\pi\Sigma, \bar{K}N} | \varphi_d P_K \rangle, \end{aligned} \quad (15)$$

which is the matrix element of the two-body T operator between the initial and final state. It contains two-body dynamics through the T operator and the kinematical input: the transformation between the Jacobi-coordinates and the deuteron wave function. This is basically the formula, which EAY used to calculate the transition amplitude from the K^-d atomic state to the $\pi\Sigma n$ continuum. As for the non-resonant amplitude they suggest to replace $T_{\pi\Sigma, \bar{K}N}$ in (15) by $V_{\pi\Sigma, \bar{K}N}$:

$$A_{BU}^{Born}(x_{\pi\Sigma}, y_N; \sigma_{\pi\Sigma}) = \langle x_{\pi\Sigma}, y_N | V_{\pi\Sigma, \bar{K}N} | \varphi_d P_K \rangle, \quad (16)$$

that is, to use the Born-approximation, which contains all the kinematics.

Thus we have three amplitudes with the properties

- $A_{BU} \longrightarrow$ three-body dynamics + three-body kinematics,
- $A_{BU}^{sing} \longrightarrow$ two-body dynamics + three-body kinematics,
- $A_{BU}^{Born} \longrightarrow$ three-body kinematics,

and we expect, that the DEV spectra

$$P_{DEV}(E_n) = P(E_n)/P^{Born}(E_n) ; \quad P_{DEV}^{sing}(E_n) = P^{sing}(E_n)/P^{Born}(E_n) \quad (17)$$

will display (reveal) three- and two-body dynamics, respectively.

It is assumed, that the E_n dependence of $P_{nonres}(E_n) = P^{Born}(E_n)$ is basically determined by the features of the initial and final states, while the details of the $V_{\bar{K}N, \pi\Sigma}$ potential (within reasonable limits) influence it only weakly. This expectation is important, if the deviation spectrum method is

to be applied for extracting some information on $\Lambda(1405)$ from an experimentally measured neutron spectrum. (It is assumed, that the matrix element (16) can be calculated in an experimental group, too.) To check this anticipated model-independence of the method, we calculated $P^{Born}(E_n)$ *not* with our realistic $\bar{K}N - \pi\Sigma$ interactions, but with the simplest possible separable potential:

$$\langle x_{\pi\Sigma} | V_{Born}^I | x_{\bar{K}N} \rangle = \frac{1}{x_{\pi\Sigma}^2 + (\beta_{\pi\Sigma}^I)^2} \lambda_{\pi\Sigma, \bar{K}N}^I \frac{1}{x_{\bar{K}N}^2 + (\beta_{\bar{K}N}^I)^2}$$

and took

$$\lambda_{\pi\Sigma, \bar{K}N}^{I=0} = \lambda_{\pi\Sigma, \bar{K}N}^{I=1} = 1, \beta_{\pi\Sigma}^{I=0} = \beta_{\pi\Sigma}^{I=1} = \beta_{\bar{K}N}^{I=0} = \beta_{\bar{K}N}^{I=1} = \beta_{Born}$$

We calculated the $P_{DEV}(E_n)$ deviation spectra for different β_{Born} values and incident kaon energies. Typical results are shown in Fig. 4. The shape of




Figure 4: Effect of β_{Born} on the deviation spectra

the P_{DEV} does not depend on the normalization of P_{nonres} (see the arbitrary choice of the $\lambda_{\pi\Sigma, \bar{K}N}$ -s) and we have normalized the P_{DEV} curves to have their maxima at 1. It can be seen, that for reasonable $\pi\Sigma - \bar{K}N$ interaction range values the spectra show practically no dependence on β_{Born} , thus confirming the presumed (approximate) model-independence of the P_{DEV}

method. According to the foresaid, the rest of our calculations were done with $\beta_{Born} = 3 fm^{-1}$.

Figs. 1., 2. and 3. demonstrate our main results. Apart from the direct spectra $P(E_n)$ - which show no trace of the $\Lambda(1405)$ - we show also the full- and single-scattering deviation spectra, too. For comparison, we plotted also the (hypothetical) $\pi^0\Sigma^0$ elastic cross sections as calculated from the $\bar{K}N - \pi\Sigma$ interaction for the corresponding $\pi\Sigma$ energy (top scale). Since this cross section "feels" only the $I = 0$ part of the $\bar{K}N - \pi\Sigma$ interaction, its peak is usually identified with the $\Lambda(1405)$ resonance. Thus the similarity of the $P_{DEV}(E_n)$ and $\sigma(\pi^0\Sigma^0)$ line shapes can tell us about the reliability of extracting information about $\Lambda(1405)$ from the reaction under consideration. Obviously, the P_{DEV}^{sing} spectra show more similarity with the original $\Lambda(1405)$ shape than the full $P_{DEV}(E_n)$ spectra, since their dynamical content is more or less the same. For three of the four considered potentials (KEK 1, KEK 2, SIDDHARTA 2) a clear resonant structure can be seen in the full deviation spectra, however, the shapes and positions can significantly differ from their "originals". The question of quantitative derivation of the $\Lambda(1405)$ resonance parameters from the (measured or calculated) deviation spectra is a separate and probably non-trivial problem and will not be discussed here.

As for the fourth potential, SIDDHARTA 1, its deviation spectra do not show any signature of its original $\Lambda(1405)$, although there are clean resonance peaks in the corresponding P_{DEV}^{sing} and $\sigma(\pi^0\Sigma^0)$ curves. The reason might be the extreme closeness of the pole to the $\bar{K}N$ threshold combined with its large width.

Finally, we asked the question, under which conditions could the resonance be observed in the direct, $P(E_n)$ spectra. For this purpose we modified two of the $\bar{K}N - \pi\Sigma$ interaction parameters of one of our potentials (KEK 1), $\lambda_{\bar{K}N, \bar{K}N}^{I=0}$ and $\lambda_{\pi\Sigma, \bar{K}N}^{I=0}$, in such a way, that the position of the $\Lambda(1405)$ remained at its original place, while its width could be made smaller. The results are shown in Fig.5.

It is seen, that in order to show up in the direct $P(E_n)$ spectra the width of the original resonance should not exceed $10 - 15 MeV$, while the deviation spectra reproduce the original resonance shape in an acceptable way. Thus the real $\Lambda(1405)$ peak, which in all models has a width of $\sim 50 - 100 MeV$ has little chance to be seen directly in this reaction, at least in the considered energy region.




Figure 5: Effect of decreasing $\Gamma_{\Lambda(1405)}$ on the neutron spectra

5. Conclusions

Dynamically exact Faddeev-type calculations for the $K^- + d \rightarrow \pi + \Sigma + n$ reaction were performed in the energy range $E_{K^-}^{cm} = 0 - 50 \text{ MeV}$ in order to find the signature of the $\Lambda(1405)$ resonance in the observable neutron spectra. Four different phenomenological $\bar{K}N - \pi\Sigma$ interactions were used, all well reproducing the experimental data in the two-body sector, but having rather different pole structure. It was shown, that due to strong kinematical masking effect the inclusive neutron spectra do not exhibit a peak, corresponding to the $\Lambda(1405)$ resonance. We demonstrated, that the deviation spectrum method in most of the cases can eliminate the disturbing kinematical factors and produce "deviation" spectra clearly showing a resonance peak, which is

correlated with the "original" two-body $\Lambda(1405)$ resonance. However, the shape and position of the peaks in the deviation spectra may significantly differ from those of their "original" counterparts. For one of the potentials (SIDDHARTA 1) even the deviation spectra show no trace of the resonance, probably due to the closeness of the original pole to the $\bar{K}N$ threshold and its large imaginary part.

Acknowledgment

The work was supported by the OTKA grant T71989.

References

- [1] J.Esmaili,Y. Akaishi and T. Yamazaki, Phys. Rev. C **83**, 055207, (2011).
- [2] K. Miyagawa and J. Haidenbauer, arXiv:1202.4272 [nucl-th] (2012).
- [3] N. V. Shevchenko, A. Gal and J. Mareš, Phys. Rev. Lett. **98** 082301 (2007), N. V. Shevchenko, A. Gal, J. Mareš and J. Révai, Phys. Rev. C **76**, 044004 (2007).
- [4] Y. Ikeda and T. Sato, Phys. Rev. C **76**, 035203 (2007).
- [5] N. V. Shevchenko, Phys. Rev. C **85**,034001 (2012).
- [6] N. V. Shevchenko, arXiv:1201.3173 [nucl-th], (2012)
- [7] A. Bahaoui, C. Fayard, T. Mizutani, and B. Saghai, Phys. Rev. C **68**, 064001 (2003).
- [8] G. Toker, A. Gal, and J. M. Eisenberg, Nucl. Phys. **A 362**, 405 (1981).
- [9] M.Torres, R.H. Dalitz and A. Deloff, Phys. Lett. **B 174**,213,(1986)
- [10] T. H. Tan, Phys. Rev. **D 7**,600,(1973)
- [11] D. Jido, E. Oset and T. Sekihara, Eur. Phys. J. A **42**, 257 (2009).
- [12] D. Jido, E. Oset and T. Sekihara, Eur. Phys. J. A **47**, 42 (2011).
- [13] A. Alaylioglu, D. Eyre, M. Brannigan and J. P. Svenne, J. Comp. Phys. **62**, 383 (1986).